

Review Article

Effect of Aerobic Training on Homocysteine and Lipid Profiles in Adults: A Systematic Review and Meta-Analysis



Omid Zafarmand¹, Fatemeh Kazeminasab^{2*}

1. Master of Exercise Physiology, Department of Physical Education and Sport Science, School of Humanities, University of Yasouj, Yasouj, Iran.
2. Assistant Professor, Department of Physical Education and Sport Science, School of Humanities, University of Kashan, Kashan, Iran.



Citation Zafarmand O, Kazeminasab F. [Effect of Aerobic Training on Homocysteine and Lipid Profiles in Adults: A Systematic Review and Meta-Analysis]. *Internal Medicine Today*. 2023; 29(4): 191-205

:<https://doi.org/10.22034/imti.2023.29.4.191>

ABSTRACT



Received: 15 Nov 2023

Accepted: 08 Jun 2024

Available Online: 12 Oct 2024

Key words:

Aerobic exercise,
Cardiovascular
diseases,
Homocysteine,
Lipid profile,

Aims A sedentary lifestyle, which is a known risk factor for several diseases, including cardiovascular disorders, is associated with elevated homocysteine levels. Studies have demonstrated that physical training can influence homocysteine levels in the body. The present study aimed to investigate the effect of aerobic exercise on homocysteine and lipid profiles in adults.

Materials & Methods A systematic search of English and Persian articles published in PubMed, Web of Science (WOS), Scopus, Scientific Information Database (SID), Noor Mags, and Magiran databases was conducted up to June 2024. The mean difference and 95% confidence intervals (CIs) were calculated using a random-effects model. Heterogeneity was assessed using the I^2 statistic, and publication bias was evaluated through visual inspection of the funnel plot and Egger's test.

Findings A total of 19 studies comprising 511 adults were included in this meta-analysis. The results indicated that aerobic exercise significantly reduced homocysteine levels [WMD= -0.915, 95% CI: -1.427 to -0.403, $P= 0.001$], LDL [WMD= -8.934 mg/dL, 95% CI: -13.555 to -4.314, $P= 0.001$] and total cholesterol (TC) [WMD= -7.390 mg/dL, 95% CI: -13.881 to -0.900, $P= 0.026$] compared to control groups. However, no significant changes were observed in triglycerides (TG) [WMD= -3.418 mg/dL, 95% CI: -9.466 to 2.611, $P= 0.266$] and high-density lipoprotein (HDL) [WMD= 1.940 mg/dL, 95% CI: -0.920 to 4.799, $P= 0.148$].

Conclusion This meta-analysis suggests that aerobic exercise in adults is associated with reduced levels of homocysteine, total cholesterol, and low-density lipoprotein (LDL). Therefore, it can be recommended as a practical and non-pharmacological strategy to help prevent cardiovascular disease and diabetes.

Corresponding Author:

Fatemeh Kazeminasab

Address: Faculty of Humanities, University of Kashan, Ghotb-e-Ravandi Blvd, Kashan, Iran.

Tel: +983155913706

Email: fkazeminasab@kashanu.ac.ir



مقاله مروری

تاثیر تمرینات هوازی بر هموسیستئین و پروفایل لیپیدی در بزرگسالان: مرور نظام مند و فراتحلیل

امید ظفرمند^۱، فاطمه کاظمی نسب^{۲*}

۱. کارشناسی ارشد فیزیولوژی ورزشی، گروه تربیت بدنی و علوم ورزشی، دانشکده علوم انسانی، دانشگاه یاسوج، یاسوج، ایران.
۲. استادیار، گروه تربیت بدنی و علوم ورزشی، دانشکده علوم انسانی، دانشگاه کاشان، کاشان، ایران.

Use your device to scan and read the article online



Citation Zafarmand O, Kazeminasab F. [Effect of Aerobic Training on Homocysteine and Lipid Profiles in Adults: A Systematic Review and Meta-Analysis]. *Internal Medicine Today*. 2023; 29(4): 191-205



<https://doi.org/10.22034/imtj.2023.29.4.191>

چکیده

تاریخ دریافت: ۱۴۰۲/۰۸/۲۵

تاریخ پذیرش: ۱۴۰۳/۰۳/۱۸

تاریخ انتشار: ۱۴۰۳/۰۷/۲۰

هدف سبک زندگی کم تحرک با افزایش سطح هموسیستئین در ارتباط است که یکی از عوامل خطر شناخته شده برای بسیاری از بیماری ها، از جمله بیماری های قلبی عروقی محسوب می شود. مطالعات نشان داده اند که فعالیت بدنی می تواند بر سطح هموسیستئین در بدن تأثیرگذار باشد. هدف از این مطالعه، بررسی اثر تمرینات هوازی بر سطح هموسیستئین و پروفایل لیپیدی در بزرگسالان بود.

مواد و روش ها: جستجوی سیستماتیک مقالات فارسی و انگلیسی منتشر شده در پایگاه های اطلاعاتی PubMed، Web of Science، SID، Scopus، Magiran و Noor Mags تا ژوئن ۲۰۲۴ انجام شد. اختلاف میانگین و فاصله اطمینان ۹۵٪ با استفاده از مدل اثرات تصادفی محاسبه گردید. ناهمگنی مطالعات با استفاده از آزمون آماری I^2 بررسی شد و سوگیری انتشار با تحلیل بصری نمودار فونل پلات و آزمون Egger ارزیابی گردید.

یافته ها: در مجموع ۱۹ مطالعه و ۵۱۱ آزمودنی بزرگسال وارد فراتحلیل حاضر شدند. نتایج نشان داد که تمرین هوازی سبب کاهش معنادار هموسیستئین $[WMD=-0.915 (-0.403 \text{ to } -1.427), P=0.001]$ ، LDL $[WMD=-0.390 (-0.900 \text{ to } 0.120), P=0.001]$ ، TC $[WMD=-0.893 (-1.381 \text{ to } -0.405), P=0.001]$ و TG $[WMD=-0.266 (-0.446 \text{ to } -0.086), P=0.001]$ را در بزرگسالان نشان داد. اما تمرین هوازی سبب تغییر معنادار HDL $[WMD=0.184 (0.018 \text{ to } 0.350), P=0.001]$ و $[WMD=1.940 (1.799 \text{ to } 2.081), P=0.001]$ نسبت به گروه کنترل در بزرگسالان نشد.

نتیجه گیری: نتایج فراتحلیل حاضر نشان داد که تمرین هوازی در بزرگسالان سبب کاهش معنادار هموسیستئین، TC و LDL سرمی می شود و بنابراین می تواند به عنوان یک راهکار غیردارویی و کاربردی برای کاهش بیماری های قلبی - عروقی و دیابت به افراد پیشنهاد شود.

کلیدواژه ها:

تمرین هوازی
 هموسیستئین
 پروفایل لیپیدی
 بیماری های قلبی عروقی

نویسنده مسئول:

فاطمه کاظمی نسب

نشانی: دانشکده علوم انسانی، دانشگاه کاشان، بلوار قطب راوندی، کاشان، ایران

تلفن: +۹۸۳۱۵۵۹۱۳۷۰۶

پست الکترونیکی: fkazeminasab@kashanu.ac.ir

Introduction

Obesity is caused by environmental and genetic factors and can be considered a serious condition associated with cardiovascular problems [1]. Nowadays, cardiovascular disease is recognized as one of the leading causes of death in the world, and many factors are involved in the development of this disease. The most important risk factors for these diseases include insulin resistance, high waist-to-hip ratio [2], low-density lipoprotein (LDL) and very low-density lipoprotein (VLDL) [3] and high-density lipoprotein (HDL) [4], total cholesterol (TC), triglycerides (TG) [5], lipid disorders and their oxidation, inappropriate dietary concentrations, inactivity, obesity, smoking, high blood pressure, and increased homocysteine levels [3, 4, 6].

Homocysteine is an amino acid in the blood that is formed from the demethylation of methionine and is known as a homologue of cysteine. Epidemiological studies have indicated that high levels of homocysteine in blood plasma are a risk factor for coronary artery disease, heart attack, and peripheral vascular disease, and cause atherosclerosis in three ways: 1) damage to the inner wall of the arteries, 2) interference with the work of blood clotting factors, 3) oxidation of LDL [7]. The reduction of homocysteine levels decreases the risk of heart attacks and strokes [8]. Plasma homocysteine levels increase with age, which has been shown to be more common in women [9]. Factors contributing to higher total homocysteine concentrations in adults include low serum folate concentrations and an increased prevalence of vitamin B12 deficiency as a result of malabsorption of these vitamins from the intestines in adults [10]. Many studies have revealed that the association between homocysteine levels and atherosclerosis is even stronger than the association between atherosclerosis and cholesterol. On the other hand, increased homocysteine concentrations are themselves an independent risk factor for coronary artery disease [11].

Nowadays, the positive effects of exercise and physical activity for the primary and secondary prevention of cardiovascular diseases have been well established. Exercise and physical activity are generally associated with a healthy lifestyle [12]. Physical activity improves homocysteine and several biochemical variables that can affect the homocysteine metabolism pathway. Oxidative stress may play a more critical role in this regard. Several studies investigating the effect of physical activity on cardiovascular risk factors support the use of aerobic

activities, such as light jogging, mountain climbing, long-distance walking, and swimming. The results of these studies show that aerobic exercise reduces plasma homocysteine levels [13, 14]. In a meta-analysis, researchers examined the effect of aerobic exercise alone in 22 studies involving 520 subjects and reported that aerobic exercise increased plasma homocysteine levels [15]. On the other hand, Silva et al. (2020) reviewed a systematic review of 22 studies and reported that exercise training increases plasma homocysteine levels [16].

Elevated homocysteine levels, especially in individuals with a sedentary lifestyle, are known to be a risk factor for chronic diseases, including cardiovascular disease, type 2 diabetes, and cognitive impairment. Moreover, lipid profile abnormalities, such as increased LDL and TG, and decreased HDL, are directly associated with the risk of atherosclerosis and metabolic syndrome. Studies have shown that physical activity, particularly aerobic exercise, can play a significant role in modulating these indicators [17].

Several physiological mechanisms have been proposed to explain the beneficial effects of aerobic exercise on blood homocysteine and lipid levels. In the context of homocysteine, regular aerobic exercise may reduce homocysteine levels by reducing oxidative stress, increasing antioxidant capacity, expanding the remethylation of homocysteine to methionine through pathways dependent on vitamins B6, B12, and folate, and increasing the metabolic utilization of methionine [15]. On the other hand, aerobic exercise can reduce TG, LDL, and TC and increase HDL by stimulating the secretion of epinephrine and norepinephrine, activating lipoprotein lipase, increasing enzymes involved in lipid metabolism such as LCAT, and improving reverse cholesterol transport. In addition, increasing the utilization of fatty acids by muscles during regular exercise is another pathway modulating plasma lipids [18].

Given the conflicting findings in previous studies and differences in design, type of exercise, intensity, and duration of intervention, a meta-analysis seems necessary to investigate the effect of aerobic exercise on homocysteine levels and lipid profiles in adults. The present study aimed to systematically analyze the available evidence and provide an overall estimate of the effectiveness of aerobic exercise on these metabolic indicators in adults.

Materials and Methods

The present study is a systematic review and meta-analysis conducted according to the Cochrane and Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [19, 20].

Data Sources and Search Methods

A search was conducted in databases to retrieve relevant research articles. To this end, Scopus, PubMed, and Web of Science (WOS) databases were searched for English-language articles up to June 2024 (without limiting the year of publication) using the following keywords:

"Aerobic Exercise", "HIIT", "Interval training", "Aerobic Interval Training", "Aerobic Interval", "Intermittent Training", "High Intensity Intermittent", "High Intensity Interval Exercise", "High Intensity", "High Intensity Interval Training", "Sprint Interval Exercise", "SIT", "Moderate-Intensity Continuous Exercise", "Homocysteine", "Lipid profile", "serum lipids", "Lipid profile", "Adults".

In addition, a search was conducted in the Scientific Information Database (SID) and Magiran databases for Persian articles up to June 2024 using the following keywords (in Persian):

"aerobic exercise", "interval exercise", "aerobic interval training", "interval training", "high-intensity interval training", "sprint interval training", "moderate-intensity interval training", "moderate-intensity continuous exercise", "homocysteine", "lipid profile", "serum lipids", "lipid profile", and "adults."

Then, the reference lists of the retrieved articles and the articles citing them were manually reviewed in Google Scholar. The database searches were conducted independently by two researchers.

Inclusion and Exclusion Criteria

To conduct the meta-analysis, articles with the following characteristics were included in the study: 1) Randomized clinical trials (RCT) and non-randomized studies (NRS), published in Persian or English; 2) studies conducted on adults; 3) studies investigating the effect of aerobic exercise versus a control group; 4) studies measuring homocysteine and lipid profile (TC, TG, HDL, LDL); 5) those studies that included mean and standard deviation data for the post-test and pre-test of the variables mentioned above for subjects of both groups (aerobic exercise and control). Exclusion criteria involved animal studies, abstracts of articles presented at conferences, theses, cross-sectional studies, reviews, and meta-analyses. The initial evaluation of the articles was conducted independently by both researchers.

Data Extraction

Information on study type, first author, year of publication, randomization or non-randomization, sample size, subject characteristics including age, gender, and exercise protocol (type of intervention, length of intervention, number of sessions per week,

and exercise intensity) was extracted [21-24]. If there was insufficient data to perform the meta-analysis, the corresponding author was contacted via email, and the data required for the present meta-analysis were obtained. Moreover, if the corresponding author did not respond or did not receive the article, data were extracted from the article chart using the Get Data software [21-24].

Quality Assessment of Articles

The quality of studies was assessed using Pedro's 9-question checklist [21-24]. The evaluation criteria included the following: 1) clarity of the eligibility criteria for subjects, 2) random assignment of participants to different groups, 3) participants to be not familiar with their groupings, 4) similarity of the subjects in terms of body weight in different study groups, 5) blind assessment for the primary research variable (blinding of all assessors), 6) dropping out of less than 15% of participants of the study, 7) intention to treat analysis, 8) reporting of statistical differences between groups for the primary research variable, 9) reporting of the mean, standard deviation, and significance level (P-value). All questions on Pedro's checklist were answered with two options: yes ✓ or no ×. The minimum score was zero, and the maximum was 9, where a higher numerical value indicated a higher quality of the study.


Meta-analysis

The present meta-analysis study was conducted to compare the effect of aerobic exercise on homocysteine and lipid profiles in adults. In this study, mean, standard deviation, and sample size were employed for statistical analysis. Data analysis was calculated using a random-effects model, and the mean difference and 95% confidence interval were considered. To determine the heterogeneity of studies, the (I^2) test was used; according to the Cochrane guidelines, heterogeneity was interpreted as less than 25% = mild heterogeneity, 25-50% = low heterogeneity, 50-75% = moderate heterogeneity, and higher than 75% = high heterogeneity [21-24]. It is worth noting that a random-effects model was employed to analyze the data. In the absence of heterogeneity, sensitivity analysis was performed using the leave-one-out method with (I^2) less than 25 as the criterion. Publication bias was also assessed using a visual interpretation of the Funnel Plot. If bias was observed, Egger's test was used as a secondary deterministic test, with a significance level of 0.1 considered significant for publication bias [25]. Statistical analysis tests were performed using CMA (version 2.0) software [25].

Results

Based on a search in scientific databases up to June 2024, 182 articles were found. After removing 85 duplicate articles and screening the titles and abstracts of the remaining articles, 43 were finally selected for full-text evaluation. After screening the full text of the articles, 24 articles were excluded from the present

meta-analysis due to the lack of post-test data, insufficient research variables, or the absence of a control group [43, 44]. Finally, 19 studies were included in the present meta-analysis. Therefore, there were 19, 7, 7, 7, and 7 studies for the homocysteine, TG, TC, HDL, and LDL variables, in respective order.



Internal Medicine Today

Figure 1. Study selection flowchart

Table 1. Characteristics of subjects and exercise protocol

Exercise intensity (%)	Workout duration (min)	Length of intervention in weeks (Number of sessions per week)	Body mass index (kg/m)	Age (Year)	Variables	Characteristics of the subjects	Sample (Gender)	Study Type-Country	Ref.
Running training at an intensity of 50% to 100% of maximum aerobic speed	-	8 (3)	Exercise: 29.50±4.50 Control: 31.40±2.60	HIIT Exercise (14 people): 24.20±4.80 Control (14 people): 22.90±5.30	Homocysteine TC TG HDL LDL	Polycystic ovary syndrome	28 females	RCT – Iran	Mohammadi et al. 2023 [26]
Continuous running training at an intensity of 65% to 75% of maximum heart rate	20 to 35	8 (3)	Exercise: 30.20±3.90 Control: 28.80±4.51	Exercise (15 people): 46.60±5.83 Control (15 people): 45.50±8.12	Homocysteine	Non-athlete	30 females	RCT – Iran	Porsesh et al. 2022 [27]
Treadmill running	20 to 35	8 (3)	HIIT training:	HIIT Exercise (11 people):	Homocysteine	Cardiovascular patients	32 males and	RCT – Iran	Avazpour et al. 2021

Exercise intensity (%)	Workout duration (min)	Length of intervention in weeks (Number of sessions per week)	Body mass index (kg/m)	Age (Year)	Variables	Characteristics of the subjects	Sample (Gender)	Study Type-Country	Ref.
training at an intensity of 70% to 100% VO ₂ peak with an activity-to-rest ratio of 15 s to 1 min			28.30±1.40 Control: 28.70±2.10	54.60±3.70 Control (11 people): 57.80±3.40			females		[28]
Walking exercise at an intensity of 60% to 70% of maximum heart rate	75	16 (2)	Exercise: 27.21±4.32 Control: 28.74±4.40	Exercise (15 people): 69.68±11.30 Control (18 people): 67.61±6.00	Homocysteine TC TG HDL LDL	Type 2 diabetes	33 females	RCT – Brazil	Silva et al 2020 [29]
Exercise bike workout: Ten 1-minute intervals at an intensity of 85% to 90% of maximum heart rate and 1 min recovery at an intensity of 50% to 55% of maximum heart rate	20	8 (3)	Exercise: 32.10±1.93 Control: 31.42±2.61	Exercise (15 people): 23.94±3.14 Control (15 people): 23.18±2.36	Homocysteine HDL LDL	Overweight and obese	30 males	RCT – Iran	Mohammadyari et al 2018 [13]
Aerobic exercise at an intensity of 65% to 75% of maximum heart rate	55 to 60	8 (3)	Exercise: 31.00±5.98 Control: 31.65±5.75	Exercise (20 people): 32.00±1.05 Control (20 people): 31.00±8.67	Homocysteine TC TG HDL LDL	Obese	40 females	RCT-Iran	Tofighi et al. 2017 [30]
Pedalling exercise with an electric bike at an intensity of 40% to 70% of maximum heart rate reserve	30	8 (3)	Exercise: 21.00±2.81 Control: 23.04±1.79	Exercise (8 people): 54.75±9.32 Control (8 people): 60.75±9.91	Homocysteine	Hemodialysis patients	16 females and males (11 females and 5 males)	RCT-Iran	Karimi et al. 2017 [14]
Running training at an intensity of 50% to 70% of maximum oxygen consumption	20 to 38	10 (3)	Exercise: 23.16±4.17 Control: 22.16±2.28	Exercise (13 people): 21.41±2.15 Control (13 people): 20.66±1.82	Homocysteine	Sedentary and obese	26 males	RCT-Iran	Aghamohammadi et al. 2015 [31]
Running or walking on a treadmill at 90% to 95% of maximum oxygen consumption	10 to 45	8 (3)	Exercise: 26.10±6.50 Control: 26.50±5.00	HIIT Exercise (10 people): 27.20±5.50 Control (10 people): 27.20±5.50	Homocysteine TC TG HDL LDL	Polycystic ovary syndrome	20 females	RCT – Norway	Almenning et al. 2015 [32]
Running training at an intensity of 60% to 75% of maximum oxygen consumption	60	8 (3)	Exercise: 25.59±1.95 Control: 26.60±1.95	Exercise (15 people): 41.46±3.10 Control (15 people): 41.06±4.00	Homocysteine	Type 2 diabetes	30 females	RCT – Iran	Hejazi et al. 2013 [33]
Bruce test exercise: running on a treadmill at an intensity of 60% to 75% of maximum heart rate	20 to 35	8 (3)	Exercise: 23.14±3.87 Control: 22.43±3.45	Exercise (14 people): 19.07±0.92 Control (14 people): 19.21±1.53	Homocysteine	Non-athlete	28 males	RCT – Iran	Bahram et al. 2013 [34]
Continuous running training at an	45 to 60	3 months (3)	Exercise: 27.43±2.78 Control:	Exercise (11 people): 44.73±4.43	Homocysteine TC TG	Healthy and inactive	21 males	RCT – Iran	Bizheh et al. 2012 [35]

Exercise intensity (%)	Workout duration (min)	Length of intervention in weeks (Number of sessions per week)	Body mass index (kg/m)	Age (Year)	Variables	Characteristics of the subjects	Sample (Gender)	Study Type-Country	Ref.
intensity of 75% to 85% of maximum heart rate			27.58±3.22	Control (10 people): 41.16±8.03					
Treadmill running exercise at an intensity of 60% to 70% of maximum oxygen consumption	20 to 40	6 (2)	Practice: - Control: -	Exercise (9 people): 00.68±00.7 Control (9 people): 00.68±00.8	Homocysteine	Parkinson's disease	18 females and males (8 females and 10 males)	RCT – New York	Donoghue et al. 2012 [36]
Treadmill running or intensity training at 60% to 75% of maximum heart rate	30 to 45	12 (3)	Exercise: 34.94±1.49 Control: 35.59±2.47	Exercise (10 people): 20 to 30 years old Control (10 people): 20 to 30 years old	Homocysteine	Obese	20 females	RCT – Iran	Taghian et al. 2011 [37]
Treadmill running training at an intensity of 50% to 70% of heart rate reserve	20 to 60	16 (3)	Exercise: 26.29±4.16 Control: 31.25±4.39	Exercise (15 people): 60 to 75 years old Control (15 people): 60 to 75 years old	Homocysteine	Non-athlete	30 females	RCT – Iran	Taghian and Esfarjani 2011 [38]
Rhythmic and step exercises at an intensity of 65% to 75% of maximum heart rate	60	8 (4)	Exercise: 32.63±3.55 Control: 30.58±2.32	Exercise (12 people): 21.81±2.40 Control (12 people): 21.50±2.56	Homocysteine	Obese	24 females	RCT – Iran	Azarniveh et al. 2011 [39]
Walking exercise on a treadmill at an intensity of 70% to 80% of maximum heart rate	30	8 (3)	Exercise: 22.00±1.99 Control: 21.50±1.42	Training (18 people): 21.50±2.28 Control (13 people): 21.75±2.67	Homocysteine TC TG HDL LDL	Inactive	31 females and males (18 females and 13 males)	NRC – Turkey	Sabasi et al. 2009 [40]
Aerobic exercise in a standing position, increasing intensity at 40% to 65% of maximum heart rate	15 to 40	8 (3)	Exercise: 25.56±4.94 Control: 26.43±3.85	Exercise (14 people): 67.38±3.37 Control (10 people): 67.30±4.73	Homocysteine	Non-athlete	24 females	RCT – Iran	Dehghan et al. 2009 [41]
Walking exercise on a treadmill at an intensity of 70% to 80% of maximum heart rate	30	8 (3)	Exercise: 22.00±1.99 Control: 21.50±1.42	Exercise (22 people): 21.50±2.28 Control (18 people): 21.75±2.66	Homocysteine TC TG HDL LDL	Healthy	40 females and males (23 females and 17 males)	RCT- Turkey	Gelecek et al. 2006 [42]

Internal Medicine Today

Characteristics of Subjects

A total of 511 subjects were included in the present meta-analysis, all of whom were adults. In addition, 261 subjects were in the aerobic exercise group with a mean age of 42.45±4.23 years and a mean BMI of 26.77±4.27 years, while 250 subjects were in the control group with a mean age of 43.08±5.99 years and a mean BMI of 27.26±3.64 years. All participants were inactive before the start of the exercise protocol, and in all studies, the

control group did not engage in any physical activity. The number of subjects in the studies ranged from a minimum of 16 [14] to a maximum of 45 [45].

Features of the Exercise Protocol

A total of 19 studies were included in the present meta-analysis, with a minimum of 20 min [13] and a maximum of 75 min [29] for each session of aerobic exercise. The exercise intensity for aerobic exercise was

at least 40–70% [14] and at most 70–100% of maximal oxygen uptake [28].


Meta-analysis Results Homocysteine

The analysis of data from 19 interventions indicated that aerobic exercise significantly reduced homocysteine [WMD= -0.915, 95% CI: -1.427 to -0.403, $P=0.001$] compared to the control group in adults (Figure 2). Heterogeneity was examined using the (I^2) test, and the results indicated that there was

high and significant heterogeneity ($P=0.001$, $I^2=88.841$).


TC

Analysis of the data from the seven interventions showed that aerobic exercise significantly reduced TC [WMD= -7.390 mg/dL, 95% CI: -13.881 to -0.900, $P=0.026$] compared to the control group in adults (Figure 3). Heterogeneity was assessed using the (I^2) test, and the results showed no significant heterogeneity ($P=0.947$, $I^2=0.000$).



Meta Analysis

Figure 2. Forest plot of the effect of aerobic exercise on homocysteine levels in adults.



Meta Analysis

Figure 3. Forest plot of the effect of aerobic exercise on TC levels in adults.

TG

Analysis of the data from the seven interventions showed that aerobic exercise did not significantly change TG [WMD= -3.418 mg/dL, 95% CI: -9.466 to 2.611, P= 0.266] compared to the control group in adults (Figure 4). Heterogeneity was examined using the (I^2) test, and the results indicated no significant heterogeneity (P=0.909, I^2 =0.000).

HDL

Analysis of the data from the seven interventions showed that aerobic exercise did not significantly change HDL [WMD= 1.940 mg/dL, 95% CI: -0.920 to

4.799, P= 0.148] compared to the control group in adults (Figure 5). Heterogeneity was assessed using the I^2 test, and the results indicated no significant heterogeneity (P=0.300, I^2 =17.044).

LDL


Data analysis of the seven interventions showed that aerobic exercise significantly reduced LDL [WMD= -8.934 mg/dL, 95% CI: -13.555 to -4.314, P= 0.001] compared to the control group in adults (Figure 6). Heterogeneity was assessed using the I^2 test, and the results showed no significant heterogeneity (P = 0.493, I^2 =0.000).

Table 2. Assessment of the quality of studies.

Study-year	1	2	3	4	5	6	7	8	9	Score
Mohammadi et al. 2023 [26]	✓	✓	×	×	✓	✓	×	✓	✓	6
Porsesh et al. 2022 [27]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Avazpour et al. 2021 [28]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Silva et al. 2020 [29]	✓	✓	×	×	✓	✓	×	✓	✓	6
Mohammadyari et al. 2018 [13]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Tofighi et al. 2017 [30]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Karimi et al. 2017 [14]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Aghamohammadi et al. 2015 [31]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Almenning et al. 2015 [32]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Hejazi et al. 2013 [33]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Bahram et al. 2013 [34]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Bizheh et al. 2012 [35]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Donoghue et al. 2012 [36]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Taghian et al. 2011 [37]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Taghian & Esfarjani, 2011 [38]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Azarniveh et al. 2011 [39]	✓	✓	×	×	✓	✓	×	✓	✓	6
Sabasi et al. 2009 [40]	✓	✓	×	✓	✓	✓	×	✓	✓	7
Dehghan et al. 2009 [41]	✓	✓	×	×	✓	✓	×	✓	✓	6
Gelecek et al. 2006 [42]	✓	✓	×	×	✓	✓	×	✓	✓	6

Internal Medicine Today

Meta Analysis




Meta Analysis

Internal Medicine Today

Figure 4. Forest plot of the effect of aerobic exercise on TG levels in adults

Meta Analysis




Meta Analysis

Internal Medicine Today

Figure 5. Forest plot of the effect of aerobic exercise on HDL levels in adults

Meta Analysis



Meta Analysis

Internal Medicine Today

Figure 6. Forest plot of the effect of aerobic exercise on LDL levels in adults

Publication Bias

The results of Egger's test indicated no significant publication bias for homocysteine ($P = 0.363$) and HDL ($P = 0.060$), as well as non-significant results for TC ($P = 0.860$) and LDL ($P = 0.836$).

Quality of Studies

The results of the quality assessment of articles using Pedro demonstrated that the minimum and maximum quality scores were 6 and 7, respectively.

Discussion

The present meta-analysis aimed to investigate the effect of aerobic exercise on homocysteine and lipid profiles in adults. The results of 19 studies with 511 subjects indicated that aerobic exercise significantly reduced homocysteine, TC, and LDL compared to the control group in adults. On the other hand, aerobic exercise did not change substantially TG and HDL compared to the control group.

Previous studies have also reported similar results; for example, Mohammadyari et al. (2018) showed that eight weeks of low-volume, high-intensity interval training reduced plasma homocysteine levels in overweight men [13]. In addition, Karimi et al. (2017) reported a significant reduction in homocysteine in hemodialysis patients after eight weeks of aerobic training [14]. However, another study observed a non-significant decrease in homocysteine in inactive overweight middle-aged women [27]. Bahram et al.

[34] reported a non-significant reduction in homocysteine in non-athletic men [34]. These differences may be due to insufficient training volume in terms of intensity or duration, differences in fitness level, and a lack of complete control of participants' nutrition.

Possible mechanisms of homocysteine reduction after aerobic exercise include reduction of oxidative stress through increased homocysteine methylation, increased S-adenosyl methionine, and increased antioxidant capacity. In general, regular exercise training increases the metabolic requirement for methionine, which leads to a decrease in homocysteine, an intermediate in methionine metabolism [15].

Aerobic exercise can affect homocysteine levels and lipid profiles through various biological pathways. One of the essential mechanisms for reducing homocysteine after exercise is to improve the function of enzymes such as methionine synthase and increase the consumption of cofactors of vitamins B6, B12, and folate, which are involved in homocysteine metabolism and are increased by regular exercise [15]. Additionally, chronic aerobic exercise activates the AMPK pathway, leading to an increase in β -oxidation of fatty acids in mitochondria, which in turn reduces blood lipids, including triglycerides and LDL [46]. Aerobic exercise also improves the antioxidant/peroxidase ratio and inhibits the degradation of compounds, such as homocysteine, by

reducing oxidative stress. On the other hand, the increase in lipoprotein lipase in skeletal muscle and adipose tissue after exercise leads to increased triglyceride uptake and increased HDL [47]. These biochemical pathways are key mechanisms in the role of aerobic exercise as a non-pharmacological strategy for improving cardiometabolic health in adults.

In the field of lipid profile, Mohammadyari et al. (2018) reported that eight weeks of high-intensity low-volume interval training reduced LDL and increased HDL [18]. Additionally, Tofighi et al. (2017) demonstrated a significant decrease in TG, TC, and LDL, along with a substantial increase in HDL, in obese women after eight weeks of aerobic training [30]. Elming et al. (2015) also reported a decrease in TG, TC, and LDL, and an increase in HDL, in women with polycystic ovary syndrome after ten weeks of aerobic training [32]. However, Sabasi et al. (2009) reported that they did not observe a significant change in these variables in sedentary individuals after eight weeks of training [40]. Gelecek et al. (2006) reported a significant decrease in TC and LDL, and a substantial increase in HDL, in healthy individuals [42]. These differences are likely due to variations in training protocols and the physiological and pathological conditions of the participants. The mechanisms of TC, TG, and LDL reduction, and HDL increase due to aerobic exercise include increased epinephrine secretion during exercise, which, via cyclic adenosine monophosphate, increases the release of free fatty acids from adipose tissue. This process occurs through the activation of lipases [48]. Physical activity appears to increase the skeletal muscle's capacity to utilize fat instead of glycogen, leading to a decrease in blood lipid levels [49]. The enzymes involved include increased activity of lecithin cholesterol acyltransferase (LCAT), which is responsible for the transfer of esters to HDL cholesterol, and lipoprotein lipase, as well as decreased activity of cholesterol ester transfer protein (CETP), which regulates the transfer of cholesterol from HDL to other lipoproteins. These enzymatic changes, together with improved fatty acid oxidation capacity, reduce VLDL cholesterol and TG [50].

The present study has several strengths. Given that differences in exercise type, duration, or intensity can affect the overall results of the meta-analysis, the exercise protocols of the included studies in this meta-analysis varied, encompassing different aerobic exercises. The limitations of the present study include the small sample size. From the available study data, it is clear that allocation concealment was not adequately reported in all randomized trials, which

may introduce selection bias into this assessment. Additionally, none of the randomized controlled trials included in this analysis reported blinding of outcome assessors, which presents the possibility of potential selection bias in this assessment of the study. Therefore, further studies with larger sample sizes are necessary to confirm and strengthen the findings of the present study.

The results of this meta-analysis indicated that aerobic exercise can significantly reduce plasma homocysteine, total cholesterol, and serum LDL levels in adults. However, no significant effect was observed on HDL and TG levels. These findings suggest that aerobic exercise can be used as an effective and recommended non-pharmacological intervention for improving specific metabolic markers associated with cardiovascular diseases and type 2 diabetes.

However, given the heterogeneity in the type, intensity, and duration of exercise in the included studies, it is still not possible to propose a precise and general prescription for aerobic exercise that maximizes its effectiveness on lipid profiles and homocysteine levels. Therefore, future studies with more accurate designs, focusing on comparing different types of exercise protocols (in terms of intensity, duration, and kind of exercise), are necessary to determine the most effective exercise approach for the prevention and control of metabolic and cardiovascular diseases.

The present meta-analysis has significant strengths. One of the most important of these strengths is the use of data from 19 studies involving a total of 511 adult participants, which provided an adequate sample size for analyzing the effects of aerobic exercise on homocysteine and lipid profiles. Additionally, the included studies encompassed a wide range of aerobic exercise intensities and durations, thereby increasing the generalizability of the results to various types of aerobic exercise. A detailed analysis of the results, including effect size (SMD/WMD), significance level (P-value), and confidence interval range, increased the transparency and accuracy of reporting the findings. In addition, the examination of heterogeneity (I^2) and the absence of publication bias, as determined by the Egger test, further enhanced the validity of the results. Moreover, the quality of the included studies was generally assessed as moderate to good.

However, this meta-analysis also has limitations that should be taken into consideration. The significant heterogeneity observed in the homocysteine variable indicates substantial differences between studies in terms of exercise protocols, demographic characteristics, and participant physiological conditions, which could

affect the accuracy of the results. Additionally, the sample size of some included studies was relatively small, which may limit the statistical power of specific analyses. The lack of complete information in some studies on dietary control, physical fitness level, and other contributing factors is another limitation. Finally, the lack of accurate reporting of some methodological details, such as blinding of assessors or allocation concealment, could increase the possibility of bias.

Given these limitations, it is recommended that future studies be conducted with larger samples, a more systematic design, and more precise control of confounding variables to provide more definitive results and more accurate recommendations regarding the type, intensity, and duration of aerobic exercise to improve homocysteine and lipid profiles.

Ethical Considerations

Compliance with ethical guidelines

The authors declare no conflicts of interest.

References

- [1]. Santosa S, Demonty I, Lichtenstein AH, Cianflone K, Jones PJ. An investigation of hormone and lipid associations after weight loss in women. *J Am Coll Nutr.* 2007;26(3):250-8. [DOI: [10.1080/07315724.2007.10719608](https://doi.org/10.1080/07315724.2007.10719608)] [PMID]
- [2]. König D, Bissé E, Deibert P, Müller HM, Wieland H, Berg A. Influence of training volume and acute physical exercise on the homocysteine levels in endurance-trained men: interactions with plasma folate and vitamin B12. *Ann Nutr Metab.* 2003;47(3-4):114-8. [DOI: [10.1159/000070032](https://doi.org/10.1159/000070032)] [PMID]
- [3]. Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol* (1985). 2005;98(4):1154-62. [DOI: [10.1152/japplphysiol.00164.2004](https://doi.org/10.1152/japplphysiol.00164.2004)] [PMID]
- [4]. Duncan GE, Perri MG, Anton SD, Limacher MC, Martin AD, Lowenthal DT, et al. Effects of exercise on emerging and traditional cardiovascular risk factors. *Prev Med.* 2004;39(5):894-902. [DOI: [10.1016/j.ypmed.2004.03.012](https://doi.org/10.1016/j.ypmed.2004.03.012)] [PMID]
- [5]. khairandish R, Ranjbar R, Habibi A. Effects of pilates training on body composition, lipid profile and some physical Fitness Parameters in Sedentary Obese women. *Jundishapur Sci Med J.* 2018;17(1):49-61. [DOI: [10.22118/jsmj.2018.59635](https://doi.org/10.22118/jsmj.2018.59635)]
- [6]. Braith RW, Stewart KJ. Resistance exercise training: its role in the prevention of cardiovascular disease. *Circulation.* 2006;113(22):2642-50. [DOI: [10.1161/CIRCULATIONAHA.105.584060](https://doi.org/10.1161/CIRCULATIONAHA.105.584060)] [PMID]
- [7]. Gallistl S, Sudi KM, Erwa W, Aigner R, Borkenstein M. Determinants of homocysteine during weight reduction in obese children and adolescents. *Metabolism.* 2001;50(10):1220-3. [DOI: [10.1053/meta.2001.25646](https://doi.org/10.1053/meta.2001.25646)] [PMID]
- [8]. Randeve HS, Lewandowski KC, Drzewoski J, Brooke-Wavell K, O'Callaghan C, Czupryniak L, et al. Exercise decreases plasma total homocysteine in overweight young women with polycystic ovary syndrome. *J Clin Endocrinol Metab.* 2002;87(10):4496-501. [DOI: [10.1210/jc.2001-012056](https://doi.org/10.1210/jc.2001-012056)] [PMID]
- [9]. Nygård O, Refsum H, Ueland PM, Vollset SE. Major lifestyle determinants of plasma total homocysteine distribution: the hordaland homocysteine study. *Am J Clin Nutr.* 1998;67(2):263-70. [DOI: [10.1093/ajcn/67.2.263](https://doi.org/10.1093/ajcn/67.2.263)] [PMID]
- [10]. Fakhrazadeh H, Ghotbi S, Pourebrahim R, Nouri M, Heshmat R, Bandarian F, et al. Total plasma homocysteine, folate, and vitamin B12 status in healthy Iranian adults: the Tehran homocysteine survey (2003-2004)/a cross-sectional population based study. *BMC Public Health.* 2006;6:29. [DOI: [10.1186/1471-2458-6-29](https://doi.org/10.1186/1471-2458-6-29)] [PMID] [PMCID]
- [11]. Fallah S, Nouroozi V, Seifi M, Samadikuchaksaraei A, Aghdash EM. Influence of oral contraceptive pills on homocysteine and nitric oxide levels: as risk factors for cardiovascular disease. *J Clin Lab Anal.* 2012;26(2):120-3. [DOI: [10.1002/jcla.21492](https://doi.org/10.1002/jcla.21492)] [PMID] [PMCID]
- [12]. Dankner R, Chetrit A, Ken Dror G, Sela BA. Physical activity is inversely associated with total homocysteine levels, independent of C677T MTHFR genotype and plasma B vitamins. *Age (Dordr).* 2007;29(4):219-27. [DOI: [10.1007/s11357-007-9041-0](https://doi.org/10.1007/s11357-007-9041-0)] [PMID] [PMCID]
- [13]. Mohammadyari S, Abdi S, Bakhtiyari A. The effect of low volume high-intensity interval training on predictive markers of cardiovascular disease in overweight men. *Sport Physiol Manag Investig.* 2019;11(2):133-43. [Link]
- [14]. Karimi A, Azamian Jazi A, Faramarzi M, Shahidi S, Azamian Jazi Z. The effect of intermittent intradialytic aerobic exercise training on plasma homocysteine levels and physical performance in hemodialysis patients. *J Adv Med Biomed Res.* 2017;25(110):94-104. [Link]
- [15]. Deminice R, Ribeiro DF, Frajacomio FT. The effects of acute exercise and exercise training on plasma homocysteine: A meta-analysis. *PLoS One.* 2016;11(3):e0151653. [DOI: [10.1371/journal.pone.0151653](https://doi.org/10.1371/journal.pone.0151653)] [PMID] [PMCID]
- [16]. AS ES, Lacerda FV, da Mota MPG. The effect of aerobic vs. resistance training on plasma homocysteine in individuals with type 2 diabetes. *J Diabetes Metab Disord.* 2020;19(2):1003-9. [DOI: [10.1007/s40200-020-00596-z](https://doi.org/10.1007/s40200-020-00596-z)] [PMID] [PMCID]
- [17]. Solymoss BC, Bourassa MG, Campeau L, Sniderman A, Marcil M, Lespérance J, et al. Effect of increasing metabolic syndrome score on atherosclerotic risk profile and coronary artery

- disease angiographic severity. *Am J Cardiol.* 2004;93(2):159-64. [\[DOI: 10.1016/j.amjcard.2003.09.032\]](#) [\[PMID\]](#)
- [18].Ngayimbesha A, Bizimana J, Gakima M. Effect of eight weeks of exercise training on lipid profile and insulin sensitivity in obese person. *Int J Sports Exerc Med.* 2019;5(2):119-25. [\[DOI: 10.23937/2469-5718/1510119\]](#)
- [19].Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev.* 2015;4(1):1. [\[DOI: 10.1186/2046-4053-4-1\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [20].Tarsilla M. Cochrane handbook for systematic reviews of interventions. *J Multidiscipl Evalua.* 2008;6:142-8. [\[DOI: 10.56645/jmde.v6i14.284\]](#)
- [21].Zafarmand O, Moghadasi M, Mogharnasi M. The effect of aerobic training on plasma levels of leptin and adiponectin in overweight and obese children and adolescents: A systematic review and meta-analysis. *Iran J Endocrinol Metab.* 2023;25(4):351-63. [\[Link\]](#)
- [22].Zafarmand O, Mogharnasi M, Moghadasi M. The effect of exercise training on serum levels of adipokines related to energy homeostasis (adropin, asprosin) and insulin resistance in patients with type 2 diabetes or obesity: A systematic review and meta-analysis. *J Appl Health Studies Sport Physiol.* 2024;11(2):23-43. [\[DOI: 10.22049/jahssp.2024.29339.1620\]](#)
- [23].Kazemi Nesab F, Zafarmand O. Comparison of the effects of high-intensity intermittent training and moderate-intensity continuous training on cardiometabolic factors in type 2 diabetic patients: a systematic review and meta-analysis. *Feyz Med Sci J.* 2024;28(1):96-109. [\[DOI: 10.48307/FMSJ.2024.28.1.98\]](#)
- [24].Mogharnasi M, Kazeminasab F, Zafarmand O, Hassanpour N. The effect of aerobic and resistance training on omentin-1 and nesfatin-1 levels in adults: A systematic review and meta -analysis. *J Birjand Univ Med Sci.* 2024;30(4):295-315. [\[DOI: 10.61186/JBUMS.30.4.295\]](#)
- [25].Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ.* 1997;315(7109):629-34. [\[DOI: 10.1136/bmj.315.7109.629\]](#) [\[PMID\]](#)
- [26].Mohammadi S, Monazzami A, Alavimilani S. Effects of eight-week high-intensity interval training on some metabolic, hormonal and cardiovascular indices in women with PCOS: a randomized controlled trail. *BMC Sports Sci Med Rehabil.* 2023;15(1):47. [\[DOI: 10.1186/s13102-023-00653-z\]](#) [\[PMID\]](#)
- [27].Porsesh M, Ahmadi Barati S, Habibi A. The effect of aerobic activity on changes in cystatin C and homocysteine levels as cardiovascular risk factors. *Jundishapur Sci Med J.* 2022;21(3):340-51. [\[DOI: 10.32598/JSMJ.21.3.2521\]](#)
- [28].Avazpour S, Nemati J, Hemmatinafar M, Salesi M. The effect of a combined training period (aerobic and high intensity interval training) and high intensity interval training on homocysteine and ESR in cardiovascular patients. *Res Med.* 2021;45(4):40-5. [\[Link\]](#)
- [29].E Silva AS, Lacerda FV, da Mota MPG. The effect of aerobic vs. resistance training on plasma homocysteine in individuals with type 2 diabetes. *J Diabetes and Metab Disord.* 2020;19(2):1003-9. [\[DOI: 10.1007/s40200-020-00596-z\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [30].Tofighi A, Jamali B, Babaei S, Amaghani A. Effect of regular exercise on serum levels of homocysteine and lipid profile in obese female. *Med J Tabriz Uni Med Sci Health Serv.* 2017;39(5):20-7. [\[Link\]](#)
- [31].Aghamohammadi F, Mogharnasi M, Kianian T. Effects of different types of exercise training and detraining on plasma homocysteine and the factors associated with obesity in sedentary young men. *Der Pharmacia Lett.* 2015;7(9):78-83. [\[Link\]](#)
- [32].Almenning I, Rieber-Mohn A, Lundgren KM, Løvvik TS, Garnæs KK, Moholdt T. Effects of high intensity interval training and strength training on metabolic, cardiovascular and hormonal outcomes in women with polycystic ovary syndrome: A pilot study. *PLoS One.* 2015;10(9): e0138793. [\[DOI: 10.1371/journal.pone.0138793\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [33].Hejazi SM, Rashidlamir A, Jebelli A, Normematolahi S, Ghazavi SM, Soltani M. The effects of 8 weeks aerobic exercise on levels of homocysteine, HS-CRP serum and plasma fibrinogen in type II diabetic women. *Life Sci J.* 2013;10(SUPPL.1):430-5. [\[Link\]](#)
- [34].Bahram ME, Najjarian M, Sayyah M, Mojtahedi H. The effect of an eight-week aerobic exercise program on the homocysteine level and VO2max in young non-athlete men. *Feyz Med Sci J.* 2013;17(2):149-56. [\[Link\]](#)
- [35].Bizheh N, Ebrahimi Atri A, Jaafari M. The effect of three months aerobic exercise on levels of hsCRP, homocysteine, serum lipids and aerobic power in healthy and inactive middle aged men. *Daneshvar Med.* 2012;20(1):43-50. [\[Link\]](#)
- [36].DiFrancisco-Donoghue J, Lamberg EM, Rabin E, Elokda A, Fazzini E, Werner WG. Effects of exercise and B vitamins on homocysteine and glutathione in Parkinson's disease: a randomized trial. *Neurodegener Dis.* 2012;10(1-4):127-34. [\[DOI: 10.1159/000333790\]](#) [\[PMID\]](#)
- [37].Taghian F, Kargarfard M, Kelishadi R. Effects of 12 weeks aerobic training on body composition, serum homocysteine and CRP levels in obese women. *J Isfahan Med Sch.* 2011;29(149):1037-45. [\[Link\]](#)
- [38].Taghian F, Esfarjani F. Effects of aerobic training on body composition and serum homocysteine in elderly women. *Armaghan Danesh.* 2011;16(3):221-32. [\[Link\]](#)
- [39].Azarniveh M, Hosseini Kakhak SAR, Hamedinia MR, Amiriparsa T, Chamri M, Hedayati M. No effect of aerobic exercise on homocysteine plasma. *Olympic.* 2011;53(19):25-38. [\[Link\]](#)
- [40].Subasi Yesilyaprak SS, Gelecek N, Özdemir N, Örmən M. Influences of acute resistance and aerobic exercises on plasma homocysteine level and lipid profiles. *Turkish J Biochem.* 2009;34(1):9-14. [\[Link\]](#)
- [41].Dehghan S, Sharifi G, Faramarzi M. The effect of eight week low impact rhythmic aerobic training on total plasma homocysteine concentration in older non-athlete women. *Jo Mazandaran Univ Med Sci.* 2009;19(72):53-9. [\[Link\]](#)
- [42].Gelecek N, Pinar L, Özdirenc M, Bediz C, Akan P, Kozan Ö, et al. Effects of brisk walking program on plasma homocysteine level and lipid profile in sedentary young subjects. *Fizyoterapi Rehabilitasyon.* 2006;17(1):42-6. [\[Link\]](#)
- [43].Rajabi A, Akbar Nezhad Gharehlo A, Madadizadeh E, Basereh A, Khoramipoor K, Pirani H, et al. The effect of 12 weeks of aerobic exercise training with or without saffron supplementation on diabetes-specific markers and inflammation in women with type 2 diabetes: A randomized double-blind placebo-controlled trial. *Eur J Sport Sci.* 2024;24(7):899-906. [\[DOI: 10.1002/ejsc.12125\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [44].Antunes HK, De Mello MT, de Aquino Lemos V, Santos-Galduróz RF, Camargo Galdieri L, Amodeo Bueno OF, et al. Aerobic physical exercise improved the cognitive function of elderly males but did not modify their blood homocysteine levels. *Dement Geriatr Cogn Dis Extra.* 2015;5(1):13-24. [\[DOI: 10.1159/000369160\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [45].Antunes HKM, De Mello MT, De Aquino Lemos V, Santos-Galduróz RF, Galdieri LC, Bueno OFA, et al. Aerobic physical exercise improved the cognitive function of elderly males but did not modify their blood homocysteine levels. *Dement Geriatr Cogn Dis Extra.* 2015;5(1):13-24. [\[DOI: 10.1159/000369160\]](#) [\[PMID\]](#) [\[PMCID\]](#)
- [46].Zouhal H, Jacob C, Delamarche P, Gratas-Delamarche A. Catecholamines and the effects of exercise, training and

- gender. Sports Med. 2008;38:401-23. [\[DOI: 10.2165/00007256-200838050-00004\]](#) [\[PMID\]](#)
- [47]. Nikolaidis MG, Kyparos A, Spanou C, Paschalis V, Theodorou AA, Vrabas IS. Redox biology of exercise: an integrative and comparative consideration of some overlooked issues. J Exp Biol. 2012;215(10):1615-25. [\[DOI: 10.1242/jeb.067470\]](#) [\[PMID\]](#)
- [48]. Shixian Q, VanCrey B, Shi J, Kakuda Y, Jiang Y. Green tea extract thermogenesis-induced weight loss by epigallocatechin gallate inhibition of catechol-O-methyltransferase. J Med Food. 2006;9(4):451-8. [\[DOI: 10.1089/jmf.2006.9.451\]](#) [\[PMID\]](#)
- [49]. Onakpoya IJ, Heneghan CJ. Effect of the novel functional fibre, polyglycoplex (PGX), on body weight and metabolic parameters: A systematic review of randomized clinical trials. Clin Nutr. 2015;34(6):1109-14. [\[DOI: 10.1016/j.clnu.2015.01.004\]](#) [\[PMID\]](#)
- [50]. Mann S, Beedie C, Jimenez A. Differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations. Sports Med. 2014;44(2):211-21. [\[DOI: 10.1007/s40279-013-0110-5\]](#) [\[PMID\]](#) [\[PMCID\]](#)